

Non-linear non-local molecular electrodynamics with nano-optical fields

Vladimir Y. Chernyak,^{1,2,a)} Praseon Saurabh,^{3,b)} and Shaul Mukamel^{3,c)}

¹Department of Chemistry, Wayne State University, 5101 Cass Ave., Detroit, Michigan 48202, USA

²Department of Mathematics, Wayne State University, 656 W. Kirby, Detroit, Michigan 48202, USA

³Department of Chemistry, University of California, Irvine, California 92697, USA

(Received 2 September 2015; accepted 6 October 2015; published online 26 October 2015)

The interaction of optical fields sculpted on the nano-scale with matter may not be described by the dipole approximation since the fields may vary appreciably across the molecular length scale. Rather than incrementally adding higher multipoles, it is advantageous and more physically transparent to describe the optical process using non-local response functions that intrinsically include all multipoles. We present a semi-classical approach for calculating non-local response functions based on the minimal coupling Hamiltonian. The first, second, and third order response functions are expressed in terms of correlation functions of the charge and the current densities. This approach is based on the gauge invariant current rather than the polarization, and on the vector potential rather than the electric and magnetic fields. © 2015 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4934231>]

I. INTRODUCTION

Quantum optics is commonly formulated by making the semiclassical approximation whereby the fields are treated classically and matter is treated quantum mechanically.¹⁻³ The radiation-matter coupling may be described by the minimal coupling Hamiltonian where the electromagnetic fields are represented by vector potential $\mathbf{A}(\mathbf{r}, t)$ ⁴ and the matter properties enter through the current $\hat{\mathbf{j}}(\mathbf{r}, t)$ and charge density $\hat{\sigma}(\mathbf{r}, t)$ matrix elements of the desired transitions.⁵ Alternatively, the multipolar hamiltonian is used where the electromagnetic field is represented by the electric and magnetic fields and matter is expanded in electric and magnetic multipoles. In most applications, including the theory of the laser,² the lowest order of this multipolar expansion, known as the dipole (long wavelength) approximation, is sufficient to account for experimental observations.⁶

In recent years, however, there has been rapid developments,⁷⁻¹¹ both theoretical and experimental, in the field of nano-optics.¹² Notable for spectroscopy are, nanoantenna,⁷ nanoplasmonic^{8,9,13} and associated spatial and temporal resolutions of optical spectra resulting in attosecond local-field enhanced ($\times 10-10^2$) spectroscopy.^{14-17,19} This is possible because of the nano-scale field confinement (~ 10 nm),⁷ as in case of nanoantenna, and spectral bandwidth of plasmonic spectra (850–2200 THz).⁸ *Nano-optical fields* are spatially confined fields which show appreciable position (\mathbf{r})-dependence on the molecular length scale.^{12,18,20-22} Combination of a nanoantenna in tip of the probe^{12,23} and utilizing its nanoplasmonic properties has pushed the boundaries of spectral resolution of optical spectroscopy, for example, surface- and tip-enhanced absorptions and Raman spectroscopy upto single molecule level.^{10,14,15,17} Non-linear optical manipulations of

nano-particles have been reported as well.^{24,25} When the field confinement is comparable to quantum confinement of molecular orbitals, the dipole approximation may not be adequate and higher multipoles are required.^{7,26}

In this paper, we calculate matter/field energy exchange and heterodyne detected optical signals [Eqs. (15) and (49)] using non-local response functions that take all multipoles into account. Using the minimal coupling Hamiltonian for the interaction of nanoscale confined optical fields with molecules, we can fully describe the radiation by a single field (the vector potential or the electric field or the magnetic field) and we do not need to use both the electric with magnetic fields. Starting with minimal coupling Hamiltonian [Eq. (1)],^{1,27,28} we write the radiation field Hamiltonian in second quantized modes of optical field [Eq. (2)] and radiation-matter interaction Hamiltonian using gauge-invariant current density $\hat{\mathbf{J}}(\mathbf{r}, t)$ [Eq. (11)].²⁸ The material properties enter via correlation functions of the current density operator $\hat{\mathbf{j}}(\mathbf{r})$ and the charge density operators $\hat{\sigma}(\mathbf{r})$ [Eqs. (8) and (9)].

The non-local response approach is particularly suitable for describing the interaction with nano-optical fields.^{8,12} Low order multipoles are adequate when the field varies slowly on the molecular scale, which is not the case in nano-optics.⁷ Coupling the microscopic Schrödinger equation with the macroscopic Maxwell's equations^{29,31} then is simply an approximate. The non-local response provides a natural exact link for computing optical signals.

We work with gauge-dependent quantities, such as the vector $\mathbf{A}(\mathbf{r}, t)$ and the scalar potentials $A_0(\mathbf{r}, t)$, making sure that all observables are gauge invariant. Gauge-invariant systems are most naturally formulated within the Lagrange formalism that involves path integrals. The Hamiltonian formalism is a bit more tricky, since gauge invariance leads to the constraint $\nabla \cdot \mathbf{E} - 4\pi\sigma = 0$, and the gauge-invariant formalism requires restricting the space of states to the physical subspace on which the constraints are satisfied. Alternatively, one can fix the gauge by imposing a set of conditions on the potentials.

a)Electronic mail: chernyak@chem.wayne.edu

b)Electronic mail: psaurabh@uci.edu

c)Electronic mail: smukamel@uci.edu

The most suitable gauge for doing quantum calculations within the Hamilton approach is known as the Coulomb (or sometimes Hamiltonian) gauge and is defined by the condition $\nabla \cdot \mathbf{A}(\mathbf{r}, t) = 0$. In this gauge, the field variables are represented by a transverse vector potential, and the longitudinal field is represented by the Coulomb potential created by the charges and therefore becomes a material, rather than field variable, whereas the total Hamiltonian H of a system interacting with the electromagnetic field is given by^{1,28}

$$\hat{H} = \hat{H}_0 + \hat{H}_{\text{rad}} + \hat{H}_{\text{int}} = \hat{H}_m + \hat{H}_c + \hat{H}_{\text{rad}} = \hat{H}_m + \hat{H}_f, \quad (1)$$

$$\hat{H}_m = \hat{H}_0 - \hat{H}_c + \hat{H}_{\text{int}}, \quad \hat{H}_f = H_{\text{rad}} + \hat{H}_c,$$

where \hat{H}_0 is the material Hamiltonian that includes the kinetic energy of the charges, as well as the Coulomb interaction between them, \hat{H}_{rad} represents the (transverse) radiation field and can be written as Eq. (2) in second quantized form as^{28,32}

$$\hat{H}_{\text{rad}} = \sum_{q\lambda} \hbar\omega_{q\lambda} \hat{b}_{q\lambda}^\dagger \hat{b}_{q\lambda}$$

$$= \frac{1}{8\pi} \int d\mathbf{r} (\hat{\mathbf{E}}_{\text{T}}^2(\mathbf{r}) + (\nabla \times \hat{\mathbf{A}}(\mathbf{r}))^2), \quad (2)$$

with

$$[\hat{b}_{q\lambda}, \hat{b}_{q'\lambda'}^\dagger] = \delta_{q,q'} \delta_{\lambda,\lambda'} \quad (3)$$

where $b_{q\lambda}$ and $b_{q\lambda}^\dagger$ are the annihilation and creation operators, respectively, for the photon modes with wavevector \mathbf{q} and polarization λ , while $\hat{\mathbf{E}}_{\text{T}}(\mathbf{r})$ is the operator of the transverse electric field; H_{int} is the radiation-matter interaction Hamiltonian,²⁸

$$\hat{H}_{\text{int}} = - \int d\mathbf{r} \left(\hat{\mathbf{j}}(\mathbf{r}) \cdot \hat{\mathbf{A}}(\mathbf{r}) - \frac{e^2}{2mc} \hat{\sigma}(\mathbf{r}) \hat{\mathbf{A}}^2(\mathbf{r}) \right). \quad (4)$$

This can be recast as

$$\hat{H}_{\text{int}} = - \int d\mathbf{r} \hat{\mathbf{J}}_{\text{int}}(\mathbf{r}) \cdot \hat{\mathbf{A}}(\mathbf{r}), \quad (5)$$

where

$$\hat{\mathbf{J}}_{\text{int}}(\mathbf{r}) = \hat{\mathbf{j}}(\mathbf{r}) - \frac{e^2}{2mc} \hat{\mathbf{A}}(\mathbf{r}) \hat{\sigma}(\mathbf{r}) \quad (6)$$

is the *effective interaction* current. The second and the third partitions of the system Hamiltonian \hat{H} in Eq. (1) involve the matter Hamiltonian \hat{H}_m that includes the kinetic energy of the electrons and their interactions with the transverse electromagnetic field; \hat{H}_c represents the Coulomb interactions between electrons that, as is well known,³³ can be interpreted as the energy of the longitudinal electromagnetic field (in particular the longitudinal field in the Coulomb gauge is a purely material variable), so that \hat{H}_f represents the total energy of the electromagnetic field.

Gauge invariance is most conveniently formulated using the electron field creation $\hat{\psi}^\dagger(\mathbf{r})$ and annihilation $\hat{\psi}(\mathbf{r})$ operators which satisfy the Fermi commutation relations

$$\{\hat{\psi}(\mathbf{r}), \hat{\psi}^\dagger(\mathbf{r}')\} = \delta(\mathbf{r} - \mathbf{r}') \quad (7)$$

the “naive,” i.e., in the absence of the electromagnetic field, current $\hat{\mathbf{j}}(\mathbf{r})$ and charge density $\hat{\sigma}(\mathbf{r})$ operators are given by

$$\hat{\mathbf{j}}(\mathbf{r}) = \frac{e\hbar}{2mi} (\hat{\psi}^\dagger(\mathbf{r}) \nabla \hat{\psi}(\mathbf{r}) - (\nabla \hat{\psi}^\dagger(\mathbf{r})) \hat{\psi}(\mathbf{r})), \quad (8)$$

$$\hat{\sigma}(\mathbf{r}) = e \hat{\psi}^\dagger(\mathbf{r}) \hat{\psi}(\mathbf{r}). \quad (9)$$

All observables should be invariant to the following gauge transformation:

$$\hat{\psi}(\mathbf{r}) \mapsto e^{i\varphi(\mathbf{r})} \hat{\psi}(\mathbf{r}), \quad \hat{\psi}^\dagger(\mathbf{r}) \mapsto e^{-i\varphi(\mathbf{r})} \hat{\psi}^\dagger(\mathbf{r}),$$

$$\hat{\mathbf{A}}(\mathbf{r}) \mapsto \hat{\mathbf{A}}(\mathbf{r}) + \frac{e}{c} \nabla \varphi(\mathbf{r}). \quad (10)$$

A straightforward inspection shows that the following current $\hat{\mathbf{J}}(\mathbf{r})$ operator²⁸

$$\hat{\mathbf{J}}(\mathbf{r}) = \hat{\mathbf{j}}(\mathbf{r}) - \frac{e^2}{mc} \hat{\mathbf{A}}(\mathbf{r}) \hat{\sigma}(\mathbf{r}), \quad (11)$$

as well as the Hamiltonian \hat{H} [Eq. (1)], with \hat{H}_{int} defined by Eq. (5) [Eq. (11)], and the Hamiltonians \hat{H}_m , \hat{H}_c , \hat{H}_{rad} , and \hat{H}_f are preserved by the gauge transformations. $\hat{\mathbf{J}}(\mathbf{r})$ will naturally appear below when calculating the matter/field energy exchange rate.

II. THE MATTER/FIELD ENERGY EXCHANGE RATE

The optical response will be calculated via the matter-field energy exchange. To that end, we define the energy of system j as

$$W_j(t) = \text{Tr}(\hat{H}_j \varrho(t)), \quad (12)$$

with $j = m, c, \text{rad}, f$, or without a subscript, stand for the energy of matter, Coulomb (longitudinal field), transverse field, total field, and total energy, respectively. Energy conservation implies $\dot{W}(t) \equiv dW(t)/dt = 0$. At equilibrium, $\dot{\varrho}(t) = 0$ and we further have $\dot{W}_j(t) = 0$ for all j , so that there is no energy exchange between the field and the matter. When the system is driven out of equilibrium, say by external fields or currents, the field-matter energy exchange rate becomes a measure of how strongly the external field drives the system out of its equilibrium. We will refer to $\dot{W}_f = -\dot{W}_m$ as the energy exchange rate between matter and field, where the field energy includes its longitudinal component. Another approach would be to consider the energy exchange rate $\dot{W}_{\text{rad}} = -\dot{W}_m - \dot{W}_c$ between the transverse field and the system; in this second approach, the Coulomb energy, which is equal to the longitudinal field energy, is included into the system.

The energy exchange rates can be evaluated by starting with Eq. (12),

$$\dot{W}_j(t) = \text{Tr}(\hat{H}_j \dot{\varrho}(t)) = -\frac{i}{\hbar} \text{Tr}(\hat{H}_j [\hat{H}, \varrho(t)])$$

$$= -\frac{i}{\hbar} \text{Tr}([\hat{H}_j, \hat{H}] \varrho(t)) = -\frac{i}{\hbar} \langle [\hat{H}_j, \hat{H}] \rangle, \quad (13)$$

and the commutators $[\hat{H}_j, \hat{H}]$ are calculated using the electron operator commutation relations [Eq. (7)], as well as the commutation relations between the field operators.³² This gives

$$\dot{W}_f(t) = -\frac{1}{2} \int d\mathbf{r} \langle \hat{\mathbf{E}}(\mathbf{r}) \cdot \hat{\mathbf{J}}(\mathbf{r}) + \hat{\mathbf{J}}(\mathbf{r}) \cdot \hat{\mathbf{E}}(\mathbf{r}) \rangle$$

$$= -\frac{1}{2} \int d\mathbf{r} \text{Tr}((\hat{\mathbf{E}}(\mathbf{r}) \cdot \hat{\mathbf{J}}(\mathbf{r}) + \hat{\mathbf{J}}(\mathbf{r}) \cdot \hat{\mathbf{E}}(\mathbf{r})) \varrho(t)), \quad (14)$$

where $\hat{\mathbf{E}}(\mathbf{r}) = \hat{\mathbf{E}}_{\text{T}}(\mathbf{r}) + \hat{\mathbf{E}}_{\text{L}}(\mathbf{r})$, with $\hat{\mathbf{E}}_{\text{L}}(\mathbf{r})$ being the (purely material) longitudinal electric field operator that represents the Coulomb field created by the charge density. The expressions

for $\hat{W}_c(t)$ and $\hat{W}_{\text{rad}}(t)$ have a form of Eq. (14) with $\hat{E}(\mathbf{r})$ replaced by $\hat{E}_L(\mathbf{r})$ and $\hat{E}_T(\mathbf{r})$, respectively.

Note that three different current operators appear naturally in the present description: (a) $\hat{j}(\mathbf{r})$ [Eq. (8)] is the *naive* gauge-dependent current density operator and does not represent physical observable; (b) \hat{J}_{int} [Eq. (6)] is an effective interaction current operator that assures the gauge-invariance of the interaction Hamiltonian in Eq. (5); and (c) $\hat{J}(\mathbf{r})$ [Eq. (11)] is the gauge-invariant current density operator which naturally appears in Eq. (14) when the energy exchange is calculated. Since the energy exchange rate is an observable, and therefore must be expressed in terms of gauge-invariant quantities.

We next consider the field-matter energy exchange rate in a system driven by and external field/current within the semiclassical approximation which neglects spontaneous processes. This boils down to replacing in Eq. (14) the electric field $\hat{E}(\mathbf{r})$ and the vector potential $\hat{A}(\mathbf{r})$ that enters $\hat{J}(\mathbf{r})$ via Eq. (11), with their classical counterparts, represented by the external (driving) field. This results in

$$\dot{W}_f(t) = - \int d\mathbf{r} \mathbf{E}(\mathbf{r}, t) \cdot \mathbf{J}(\mathbf{r}, t), \quad (15)$$

where

$$\mathbf{J}(\mathbf{r}, t) = \langle \hat{J}(\mathbf{r}, t) \rangle = \text{Tr}(\hat{J}(\mathbf{r})\hat{\rho}(t)) \quad (16)$$

is the expectation value of the gauge-invariant current density, evaluated at the density matrix $\hat{\rho}(t)$ of the driven system. The formal arguments in support of the intuitively natural procedure, we are using in this manuscript, as well as a way of calculating the radiative corrections will be addressed elsewhere.

It follows from Eqs. (15) and (16) that the semiclassical matter-field energy exchange rate for a driven system may be expressed in terms of the expectation value of the gauge-invariant current density $\mathbf{J}(\mathbf{r}, t)$. It is thus natural to define the optical response functions as the expansion of the latter in powers of the driving field.

This can be represented in a compact form by introducing the following notation for Liouville space (tetradic) operators. With any Hilbert space operator \hat{Q} , we associate the following four Liouville space operators: *left* \hat{Q}^L , *right* \hat{Q}^R , *plus* \hat{Q}^+ , and *minus* \hat{Q}^- , defined by its action on another operator \hat{X} ,³⁴

$$\begin{aligned} \hat{Q}^L(\hat{X}) &\equiv \hat{Q}\hat{X}, & \hat{Q}^R(\hat{X}) &\equiv \hat{X}\hat{Q}, \\ \hat{Q}^+ &\equiv \frac{1}{2}(\hat{Q}^L + \hat{Q}^R), & \hat{Q}^- &\equiv \hat{Q}^L - \hat{Q}^R. \end{aligned} \quad (17)$$

Adopting the interaction picture (with time parameter τ) in Liouville space, we arrive at

$$\begin{aligned} \mathbf{J}(\mathbf{r}, t; \mathbf{A}; A_0) &= \left\langle \left(\hat{j}^+(\mathbf{r}, t) - \frac{e}{mc} \mathbf{A}(\mathbf{r}, t) \hat{\sigma}^+(\mathbf{r}, t) \right) \right. \\ &\quad \times \text{T exp} \left(-i\hbar^{-1} \int_{-\infty}^t d\tau \hat{H}_{\text{int}}^-(\mathbf{A}, A_0; \tau) \right) \Bigg\rangle, \end{aligned} \quad (18)$$

$$\begin{aligned} \hat{H}_{\text{int}}^-(\mathbf{A}, A_0; \tau) &= - \int d\mathbf{r} \left(\mathbf{A}(\mathbf{r}) \cdot \hat{\mathbf{j}}^-(\mathbf{r}, \tau) \right. \\ &\quad \left. - \frac{e}{2mc} \mathbf{A}^2(\mathbf{r}) \hat{\sigma}^-(\mathbf{r}, \tau) + A_0(\mathbf{r}) \hat{\sigma}^-(\mathbf{r}, \tau) \right), \end{aligned} \quad (19)$$

where the third term in the r.h.s. of Eq. (18) describes the interaction of the system's charge density with the longitudinal component of the driving field; in the Coulomb gauge, adopted here, it originates from the Coulomb interaction between the

system and external charge density; the latter together with the external current density is responsible for creating the external (driving) field. The interaction Hamiltonian in Eq. (5) does not have a scalar potential component, since the contribution of the scalar potential is included in the H_c of Eq. (1). The reason of explicitly including the scalar potential in Eq. (19) will be explained below.

Eq. (15) together with Eqs. (18) and (19) constitutes the main formal result of this paper. A crucial feature of $\mathbf{J}(\mathbf{r}, t; \mathbf{A}; A_0)$ [Eq. (18)] is its invariance with respect to time-dependent gauge transformations of the potentials,

$$\begin{aligned} \mathbf{A}(\mathbf{r}, t) &\mapsto \mathbf{A}(\mathbf{r}, t) + \frac{e}{c} \nabla \varphi(\mathbf{r}, t), & A_0(\mathbf{r}, t) \\ &\mapsto A_0(\mathbf{r}, t) + \frac{e}{c} \frac{\partial \varphi(\mathbf{r}, t)}{\partial t}. \end{aligned} \quad (20)$$

This property can be rationalized by noting that the transformation [Eq. (20)] in Eq. (18) is equivalent to performing the gauge transformation to the fermion fields in the interaction picture, which is given by Eq. (10), but with a time-dependent $\varphi(\mathbf{r}, \tau)$. This can be verified directly. The gauge invariance of the current $\mathbf{J}(\mathbf{r}, t; \mathbf{A}; A_0)$ then follows from the fact that the gauge transformation applied to the fermion operators does not change their commutation relations. Also, the gauge invariance of the expectation value of the gauge-invariant current is intuitively clear, since any legitimate, i.e., gauge-invariant, observable should not change upon a gauge transformation of the external (driving) field.

Gauge invariance considerably simplifies the expressions for the optical response functions, presented below. Note that Eq. (18) has been derived, in the Coulomb gauge, so that the third term in the r.h.s. of the expression for \hat{H}_{int} [Eq. (19)] is absolutely necessary. However, the gauge invariance of the optical response allows to use any gauge; for the sake of minimizing the number of terms in the expressions for the response functions, we adopt the gauge $A_0(\mathbf{r}) = 0$. Then the third term in the expression for \hat{H}_{int} should be dropped whereas the transverse vector potential should be replaced with gauge invariant potential $\mathbf{A}^{\text{inv}}(\mathbf{r}, t)$,

$$\begin{aligned} \mathbf{A}^{\text{inv}}(\mathbf{r}, t) &= \int_{-\infty}^t d\tau \mathbf{E}(\mathbf{r}, \tau) \\ &= \mathbf{A}(\mathbf{r}, t) - \int_{-\infty}^t d\tau \nabla A_0(\mathbf{r}, \tau), \end{aligned} \quad (21)$$

since, in the $A_0(\mathbf{r}) = 0$ gauge, we have $\dot{\mathbf{A}} = \mathbf{E}$. Note that the $A_0(\mathbf{r}) = 0$ gauge provides an explicitly gauge-invariant picture of the optical response, which is expressed in terms of the external electric field, or equivalently \mathbf{A}^{inv} , both are gauge invariant.

The optical response functions are defined by an expansion of the current $\mathbf{J}(\mathbf{r}, t; \mathbf{A}; A_0) = \mathbf{J}(\mathbf{r}, t; \mathbf{A}^{\text{inv}}; 0)$ in powers of n driving field \mathbf{A}^{inv} ,

$$\begin{aligned} J_k(\mathbf{r}, t) &= \sum_{k_1} \int_{-\infty}^t d\tau_1 \int d\mathbf{r}_1 \zeta_{kk_1}^{(1)}(\mathbf{r}, t; \mathbf{r}_1, \tau_1) A_{k_1}^{\text{inv}}(\mathbf{r}_1, \tau_1) \\ &\quad + \sum_{n=2}^{\infty} \frac{1}{n!} \sum_{k_1 \dots k_n} \int d\tau_1 d\mathbf{r}_1 \dots d\tau_n d\mathbf{r}_n \\ &\quad \times \zeta_{kk_1 \dots k_n}^{(n)}(\mathbf{r}, t; \mathbf{r}_1, \tau_1; \dots; \mathbf{r}_n, \tau_n) \\ &\quad \times A_{k_1}^{\text{inv}}(\mathbf{r}_1, \tau_1) \dots A_{k_n}^{\text{inv}}(\mathbf{r}_n, \tau_n), \end{aligned} \quad (22)$$

where $k, k_1, \dots, k_n = 1, 2, 3$.

The total energy exchange evaluated at time t in Eq. (15) can now be expanded order by order in driving fields. Note that time ordering is built into the response functions $\zeta^{(n)}(\mathbf{r}, t; \mathbf{r}_n, \tau_n; \dots; \mathbf{r}_1, \tau_1)$. The first order energy loss rate is calculated using Eqs. (15)–(22),

$$\Delta\dot{W}^{(1)}(t) = - \sum_{kk_1} \int d\mathbf{r} \int d\mathbf{r}_1 \int_{-\infty}^t d\tau_1 E_k(\mathbf{r}, t) A_{k_1}^{\text{inv}}(\mathbf{r}_1, \tau_1) \times \zeta_{kk_1}^{(1)}(\mathbf{r}, t; \mathbf{r}_1, \tau_1). \quad (23)$$

The first-order *non-local* response function is given by

$$\zeta_{kk_1}^{(1)}(\mathbf{r}, t; \mathbf{r}_1, \tau_1) = -i\hbar^{-1} \langle \hat{j}_k^+(\mathbf{r}, t) \hat{j}_{k_1}^-(\mathbf{r}_1, \tau_1) \rangle - \frac{e}{mc} \delta_{kk_1} \langle \hat{\sigma}^+(\mathbf{r}, t) \rangle \times \delta(t - \tau_1) \delta(\mathbf{r} - \mathbf{r}_1). \quad (24)$$

This can be recast using ordinary Hilbert space operators as

$$\zeta_{kk_1}^{(1)}(\mathbf{r}, t; \mathbf{r}_1, \tau_1) = -i\hbar^{-1} \langle [\hat{j}_k(\mathbf{r}, t), \hat{j}_{k_1}(\mathbf{r}_1, \tau_1)] \rangle - \frac{e}{mc} \delta_{kk_1} \langle \hat{\sigma}(\mathbf{r}, t) \rangle \delta(t - \tau_1) \delta(\mathbf{r} - \mathbf{r}_1). \quad (25)$$

Superoperators are a convenient book-keeping device but at the end, we can switch to ordinary operators. Now we turn to the non-linear matter/field energy exchange. Expanding

Eq. (15) to second order gives

$$\Delta\dot{W}^{(2)}(t) = - \sum_{k, k_1 k_2} \int d\mathbf{r} \int d\mathbf{r}_1 \int d\mathbf{r}_2 \int_{-\infty}^t d\tau_1 \times \int_{-\infty}^t d\tau_2 E_k(\mathbf{r}, t) A_{k_1}^{\text{inv}}(\mathbf{r}_1, \tau_1) \times A_{k_2}^{\text{inv}}(\mathbf{r}_2, \tau_2) \zeta_{kk_1 k_2}^{(2)}(\mathbf{r}t; \mathbf{r}_1 \tau_1, \mathbf{r}_2 \tau_2), \quad (26)$$

where the second order non-local response function $\zeta_{kk_2 k_1}^{(2)}(\mathbf{r}t; \mathbf{r}_2 \tau_2, \mathbf{r}_1 \tau_1)$ is

$$= -\frac{1}{2} (i\hbar^{-1})^2 \langle \hat{j}_k^+(\mathbf{r}, t) \hat{j}_{k_2}^-(\mathbf{r}_2, \tau_2) \hat{j}_{k_1}^-(\mathbf{r}_1, \tau_1) \rangle + i\hbar^{-1} \frac{e}{2mc} \left(\delta_{k_1 k_2} \langle \hat{j}_k^+(\mathbf{r}, t) \hat{\sigma}^-(\mathbf{r}_1, \tau_1) \rangle \times \delta(\mathbf{r}_2 - \mathbf{r}_1) \delta(\tau_2 - \tau_1) + 2\delta_{kk_1} \delta(\mathbf{r} - \mathbf{r}_1) \times \delta(t - \tau_1) \langle \hat{\sigma}^+(\mathbf{r}, t) \hat{\sigma}^-(\mathbf{r}_1, \tau_1) \rangle \right), \quad (27)$$

which in Hilbert space becomes

$$= -\frac{1}{2} (i\hbar^{-1})^2 \langle [[\hat{j}_k(\mathbf{r}, t), \hat{j}_{k_2}(\mathbf{r}_2, \tau_2)], \hat{j}_{k_1}(\mathbf{r}_1, \tau_1)] \rangle + i\hbar^{-1} \frac{e}{2mc} \left(\delta_{k_1 k_2} \langle [\hat{j}_k(\mathbf{r}, t), \hat{\sigma}(\mathbf{r}_1, \tau_1)] \rangle \times \delta(\mathbf{r}_2 - \mathbf{r}_1) \delta(\tau_2 - \tau_1) + 2\delta_{kk_1} \delta(\mathbf{r} - \mathbf{r}_1) \times \delta(t - \tau_1) \langle [\hat{\sigma}(\mathbf{r}, t), \hat{\sigma}(\mathbf{r}_1, \tau_1)] \rangle \right). \quad (28)$$

Finally, the third order energy exchange is given by

$$\Delta\dot{W}^{(3)}(t) = - \sum_{k, k_1 k_2 k_3} \int d\mathbf{r} \int d\mathbf{r}_1 \dots \int d\mathbf{r}_3 \int_{-\infty}^t d\tau_1 \dots \int_{-\infty}^t d\tau_3 E_k(\mathbf{r}, t) A_{k_1}^{\text{inv}}(\mathbf{r}_1, \tau_1) A_{k_2}^{\text{inv}}(\mathbf{r}_2, \tau_2) A_{k_3}^{\text{inv}}(\mathbf{r}_3, \tau_3) \zeta_{kk_3 k_2 k_1}^{(3)}(\mathbf{r}t; \mathbf{r}_3 \tau_3, \mathbf{r}_2 \tau_2, \mathbf{r}_1 \tau_1), \quad (29)$$

where the third order non-local response function, $\zeta_{kk_3 k_2 k_1}^{(3)}(\mathbf{r}t; \mathbf{r}_3 \tau_3, \mathbf{r}_2 \tau_2, \mathbf{r}_1 \tau_1)$, is given by

$$= -\frac{1}{6} (i\hbar^{-1})^3 \langle \hat{j}_k^+(\mathbf{r}, t) \hat{j}_{k_3}^-(\mathbf{r}_3, \tau_3) \hat{j}_{k_2}^-(\mathbf{r}_2, \tau_2) \hat{j}_{k_1}^-(\mathbf{r}_1, \tau_1) \rangle + (i\hbar^{-1})^2 \frac{e}{4mc} \left\{ \delta_{k_3 k_2} \langle \hat{j}_k^+(\mathbf{r}, t) \hat{j}_{k_2}^-(\mathbf{r}_2, \tau_2) \hat{\sigma}^-(\mathbf{r}_1, \tau_1) \rangle \delta(\mathbf{r}_3 - \mathbf{r}_2) \delta(\tau_3 - \tau_2) + \delta_{k_3 k_2} \langle \hat{j}_k^+(\mathbf{r}, t) \hat{\sigma}^-(\mathbf{r}_2, \tau_2) \hat{j}_{k_1}^-(\mathbf{r}_1, \tau_1) \rangle \delta(\mathbf{r}_2 - \mathbf{r}_1) \delta(\tau_2 - \tau_1) + 2\delta_{kk_1} \langle \hat{\sigma}^+(\mathbf{r}, t) \hat{j}_{k_2}^-(\mathbf{r}_2, \tau_2) \hat{j}_{k_1}^-(\mathbf{r}_1, \tau_1) \rangle \delta(\mathbf{r} - \mathbf{r}_1) \delta(t - \tau_1) \right\} - (i\hbar) \frac{e^2}{2mc^2} \delta_{kk_1} \langle \hat{\sigma}^+(\mathbf{r}, t) \hat{\sigma}^-(\mathbf{r}_1, \tau_1) \rangle \delta(\mathbf{r}_3 - \mathbf{r}_2) \delta(\mathbf{r} - \mathbf{r}_1) \delta(\tau_3 - \tau_2) \delta(t - \tau_1). \quad (30)$$

In Hilbert space, it is given by

$$= -\frac{1}{6} (i\hbar^{-1})^3 \langle [[[\hat{j}_k(\mathbf{r}, t), \hat{j}_{k_3}(\mathbf{r}_3, \tau_3)], \hat{j}_{k_2}(\mathbf{r}_2, \tau_2)], \hat{j}_{k_1}(\mathbf{r}_1, \tau_1)] \rangle + (i\hbar^{-1})^2 \frac{e}{4mc} \left\{ \delta_{k_3 k_2} \langle [[\hat{j}_k(\mathbf{r}, t), \hat{j}_{k_2}(\mathbf{r}_2, \tau_2)], \hat{\sigma}(\mathbf{r}_1, \tau_1)] \rangle \times \delta(\mathbf{r}_3 - \mathbf{r}_2) \delta(\tau_3 - \tau_2) + \delta_{k_3 k_2} \langle [[\hat{j}_k(\mathbf{r}, t), \hat{\sigma}(\mathbf{r}_2, \tau_2)], \hat{j}_{k_1}(\mathbf{r}_1, \tau_1)] \rangle \delta(\mathbf{r}_2 - \mathbf{r}_1) \delta(\tau_2 - \tau_1) + 2\delta_{kk_1} \langle [[\hat{\sigma}^+(\mathbf{r}, t), \hat{j}_{k_2}(\mathbf{r}_2, \tau_2)], \hat{j}_{k_1}(\mathbf{r}_1, \tau_1)] \rangle \delta(\mathbf{r} - \mathbf{r}_1) \delta(t - \tau_1) \right\} - (i\hbar) \frac{e^2}{2mc^2} \delta_{kk_1} \langle [[\hat{\sigma}(\mathbf{r}, t), \hat{\sigma}(\mathbf{r}_1, \tau_1)]] \rangle \delta(\mathbf{r}_3 - \mathbf{r}_2) \delta(\mathbf{r} - \mathbf{r}_1) \delta(\tau_3 - \tau_2) \delta(t - \tau_1). \quad (31)$$

Eqs. (25), (28), and (31) constitute our final exact expressions for the non-local response functions, given by correlation functions of the charge density ($\hat{\sigma}(\mathbf{r}, t)$) and the current density operator ($\hat{j}(\mathbf{r}, t)$). The current and charge densities in Eqs. (25),

(28), and (31) can be computed using the standard output of quantum chemistry packages, like Molpro,³⁵ by calculating the relevant molecular orbitals, then explicitly evaluating transition current and charge densities Eqs. (8) and (9), and,

expressing them in terms of superoperators using Eqs. (17). Note that in the non-local representation, magnetic and electric contributions need not be treated separately which greatly simplifies the analysis.

III. RECOVERING THE DIPOLE APPROXIMATION

Formally, the minimal coupling and the multipolar Hamiltonians are connected by the Power-Zienau transformation in the joint space of matter and field degrees of freedom.³⁶ However, some discrepancies arise when the two Hamiltonians are applied in practice. A known example is an extra pre-factor of $(\omega_\alpha/\Omega)^2$ with material frequency (ω_α) and field frequency (Ω) in the absorption lineshape³⁷ (see the Appendix). The issue was resolved by making the canonical transformation between the two Hamiltonians. The non-locality of the response implies that a field acting at one point includes a charge or current at a different point, suggesting an induced electronic coherence. While the multipolar expansion is computationally convenient, it hides this interesting non-local physical picture. The non-local response functions avoid the tedious canonical transformation to the multipolar Hamiltonian³⁷ and show no discrepancy between the dipole and minimal coupling absorption lineshapes.

As a consistency check, here we show how the dipole approximation for the linear response is recovered for the nonlinear minimal coupling response in agreement with the multipolar Hamiltonian. We consider a system of the size, small compared to the field wavelength, which allows us to drop the \mathbf{r} -dependence in the fields \mathbf{E} and \mathbf{A}^{inv} and recast Eq. (23) in a form

$$\Delta\dot{W}_d^{(1)} = i\hbar^{-1}\mathbf{E}(t) \cdot \mathbf{u}(t) + \frac{Ne^2}{mc}\mathbf{E}(t) \cdot \mathbf{A}^{\text{inv}}(t), \quad (32)$$

where N is the number of electrons,

$$\mathbf{u}(t) = \sum_{k,s} \int_{-\infty}^t dt' \langle \hat{u}_k^+(t) \hat{u}_s^-(t') \rangle A_s^{\text{inv}}(t'), \quad (33)$$

$k, s = 1, 2, 3$, and

$$\hat{\mathbf{u}}(t) = \int d\mathbf{r} \hat{\mathbf{j}}(\mathbf{r}, t) = - \int d\mathbf{r} \mathbf{r} (\nabla \cdot \hat{\mathbf{j}}(\mathbf{r}, t)) = \frac{d\hat{\boldsymbol{\mu}}(t)}{dt}, \quad (34)$$

with $\hat{\boldsymbol{\mu}}$ being the dipole operator. The above expressions can be interpreted as follows: the operator $\hat{\mathbf{u}}$ is the integral over the space of the naive (i.e., in the absence of the field) current density operator; the second equality in Eq. (34) is due to the Stokes formula, whereas the last equality follows from the continuity equation

$$\nabla \cdot \hat{\mathbf{j}}(\mathbf{r}, t) + \frac{d\hat{\sigma}(\mathbf{r}, t)}{dt} = 0 \quad (35)$$

in the non-driven system that holds on the operator level, combined with the well-known definition of the dipole operator $\hat{\boldsymbol{\mu}}$. As a result, $\hat{\mathbf{u}}$ can be viewed as the time derivative of the dipole. The variable $\mathbf{u}(t)$ is the expectation value of $\hat{\mathbf{u}}(t)$ in the linearly driven system; Eq. (32) is obtained by integration over \mathbf{r} and \mathbf{r}' in Eq. (23) under the assumption that the fields do not depend on \mathbf{r} .

Substituting the expression for $\hat{\mathbf{u}}(t)$ as the time derivative of the dipole [Eq. (34)] into Eq. (33), followed by substituting the latter into Eq. (32), and further integrating over t' by parts, as well as taking the derivative with respect to t out of the integral over t' , we arrive at

$$\begin{aligned} \Delta\dot{W}_d^{(1)} &= -i\hbar^{-1} \sum_{k,s} E_k(t) \frac{d}{dt} \int_{-\infty}^t dt' \langle \hat{\mu}_k^+(t) \hat{\mu}_s^-(t') \rangle A_s^{\text{inv}}(t') \\ &+ i\hbar^{-1} \sum_{k,s} E_k(t) \frac{d}{dt} \langle \langle \hat{\mu}_k^+(t) \hat{\mu}_s^-(t-0) \rangle \rangle A_s^{\text{inv}}(t) \\ &- i\hbar^{-1} \sum_{k,s} E_k(t) \langle \hat{\mu}_k^+(t) \hat{\mu}_s^-(t-0) \rangle A_s^{\text{inv}}(t) \\ &+ \frac{Ne^2}{mc} \mathbf{E}(t) \cdot \mathbf{A}^{\text{inv}}(t), \end{aligned} \quad (36)$$

where the second term in the r.h.s. of Eq. (36) is a contact (boundary) contribution in the by parts integration over t' , whereas the third term compensates for the contact (boundary contribution) that occurs in the time derivative with respect to t of an integral, whose upper limit is t -dependent. These two terms can be computed using the fact that for any two operators \hat{X} and \hat{Y} , we have $\langle \hat{X}^+(t) \hat{Y}^-(t-0) \rangle = \langle [\hat{X}(t), \hat{Y}(t)] \rangle$. This implies that the first contact term vanishes, whereas the second one is obtained using the well-known commutation relation (that can also be verified directly using a simple and straightforward calculation),

$$[\hat{j}_k(\mathbf{r}'), \hat{\sigma}(\mathbf{r})] = \frac{ie^2\hbar}{mc} \frac{\partial}{\partial r_k} \delta(\mathbf{r} - \mathbf{r}'). \quad (37)$$

After some simple and straightforward transformations, we obtain

$$[\hat{\mu}_k, \hat{u}_s] = -\frac{i\hbar Ne^2}{mc} \delta_{k,s}. \quad (38)$$

The third and fourth terms in the r.h.s. of Eq. (36) now cancel each other. Recalling the standard definition of the time-domain linear response function in the dipole approximation,

$$\chi_{k,s}^{(1)}(t) = i\hbar^{-1} \langle \hat{\mu}_k^+(t) \hat{\mu}_s^-(t) \rangle, \quad (39)$$

and the fact that $\dot{\mathbf{A}}^{\text{inv}}(t) = \mathbf{E}(t)$, we arrive at

$$\Delta\dot{W}_d^{(1)} = - \sum_{k,s} E_k(t) \frac{d}{dt} \int_{-\infty}^t dt' \chi_{k,s}^{(1)}(t-t') E_s(t') \quad (40)$$

that recovers the dipole approximation for the energy exchange rate in terms of the standard response function $\chi_{k,s}^{(1)}(t)$.

We first note that, the simple expression for the dipole approximation [Eq. (40)] was obtained by the cancellation of the last two terms in Eq. (36), which would be impossible if the second (local) term in the expression for the linear response [Eq. (23)] was omitted. The derivation presented in this section thus does not show the ‘‘extra pre-factor of $(\omega/\Omega)^2$ ’’ in absorption spectra.^{30,37} Second, if we define

$$\hat{\boldsymbol{\mu}} = \int d\mathbf{r} \hat{\mathbf{P}}(\mathbf{r}), \quad \hat{\mathbf{J}}(\mathbf{r}, t) = -\frac{d\hat{\mathbf{P}}(\mathbf{r}, t)}{dt}, \quad (41)$$

as is done in the multipolar Hamiltonian formalism with the magnetic terms neglected, one immediately obtains from

Eq. (40)

$$\Delta\dot{W}_d^{(1)} = -\mathbf{E}(t) \cdot \int d\mathbf{r} \mathbf{J}(\mathbf{r}, t), \quad (42)$$

thereby recovering the long wavelength limit of Eq. (15). Third, an attempt to compute the dipole approximation for the energy exchange rate, starting with the “standard” dipole approximation Hamiltonian,

$$\hat{H}_{\text{int}}^d = -\hat{\boldsymbol{\mu}} \cdot \mathbf{E}(t), \quad (43)$$

will result in an expression, different from Eq. (40). This can be explained as follows. The dipole Hamiltonian [Eq. (43)] is naturally obtained from the multipolar formalism that can be described as using a multipolar gauge which is different from the Coulomb gauge, or equivalently doing a canonical transformation that redistributes the energy between the matter and the field. Therefore, a calculation based on Eq. (43) addresses an exchange between energies, different from the ones, used in our calculations.

IV. HETERODYNE DETECTED SIGNALS

Heterodyne detection measures the energy exchange with a given wavevector component of the field rather than the entire field. The heterodyne detected signal can be recovered from the energy exchange rate.³⁴ We start with the (transverse) vector potential operator $\hat{\mathbf{A}}(\mathbf{r}, t)$ expanded in modes $\mathbf{q}\lambda$,

$$\hat{\mathbf{A}}(\mathbf{r}) = \sum_{\mathbf{q}\lambda} c \frac{\epsilon_{\mathbf{q}\lambda}}{\omega_{\mathbf{q}}} \mathbf{e}_{\mathbf{q}\lambda} (\hat{b}_{\mathbf{q}\lambda}^\dagger e^{-i\mathbf{q}\cdot\mathbf{r}} + \hat{b}_{\mathbf{q}\lambda} e^{i\mathbf{q}\cdot\mathbf{r}}), \quad (44)$$

where c is speed of light, $\epsilon_{\mathbf{q}\lambda} = (2\pi\hbar\omega_{\mathbf{q}}/V)^{1/2}$, and $\mathbf{e}_{\mathbf{q}\lambda}$ is the polarization vector, while V is the quantization volume. The transverse nature of the photon field manifests itself in the relation $\mathbf{q} \cdot \mathbf{e}_{\mathbf{q}\lambda} = 0$.

The heterodyne detected signal S_{het} can be defined as the rate of change of the photon number $\hat{N}_s = \hat{b}_{\mathbf{q}_s\lambda_s}^\dagger \hat{b}_{\mathbf{q}_s\lambda_s}$ in mode s ,³⁴

$$S_{\text{het}}(s) = \frac{d\langle \hat{N}_s \rangle}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{N}_s], \quad (45)$$

where the expectation value $\langle \hat{N}_s \rangle$ is taken with respect to material density matrix. More precisely, the heterodyne detected signal is the cross-component obtained by mixing the signal field (i.e., generated by the currents induced in the material by the driving field) with the heterodyne counterpart. Evaluating the commutations gives

$$S_{\text{het}}(s) = \frac{dB_{\mathbf{q}_s\lambda_s}^*}{dt} b_{\mathbf{q}_s\lambda_s} + B_{\mathbf{q}_s\lambda_s}^* \dot{b}_{\mathbf{q}_s\lambda_s} + \text{c.c.}, \quad \dot{b}_{\mathbf{q}_s\lambda_s} = \frac{db_{\mathbf{q}_s\lambda_s}}{dt}, \quad (46)$$

with

$$B_{\mathbf{q}_s\lambda_s} = \langle \hat{b}_{\mathbf{q}_s\lambda_s} \rangle_{\text{het}}, \quad b_{\mathbf{q}_s\lambda_s} = \langle \hat{b}_{\mathbf{q}_s\lambda_s} \rangle = \text{Tr}(\hat{b}_{\mathbf{q}_s\lambda_s} \hat{\rho}(t)), \quad (47)$$

so that $B_{\mathbf{q}_s\lambda_s}(t) = \bar{B}_{\mathbf{q}_s\lambda_s}(t) e^{-i\omega_{\mathbf{q}_s} t}$ is a classical field that represents the heterodyne and is linear combination of gauge-invariant electric and transverse vector potential with temporal envelopes $(\bar{E}(t))$ and $(\bar{A}(t))$ at the same mode \mathbf{q}_s , i.e., $B_{\mathbf{q}_s\lambda_s}(t) = (\bar{E}_{\mathbf{q}_s\lambda_s}(t) \pm \bar{A}_{\mathbf{q}_s\lambda_s}(t)) e^{-i\omega_{\mathbf{q}_s} t}$ where $\bar{A}_{\mathbf{q}_s\lambda_s}(t)$ is the temporal

envelop of vector potential. Whereas $b_{\mathbf{q}_s\lambda_s}(t)$ is the expectation value of the photon annihilation operator, evaluated at the density matrix $\hat{\rho}(t)$ of the driven system. The latter can be evaluated using the Heisenberg equation of motion,

$$\begin{aligned} \dot{b}_{\mathbf{q}_s\lambda_s} &= -\frac{i}{\hbar} \langle [\hat{H}, \hat{b}_{\mathbf{q}_s\lambda_s}] \rangle \\ &= -i\omega_{\mathbf{q}_s} b_{\mathbf{q}_s\lambda_s} + \frac{i}{\hbar} \int d\mathbf{r} \langle \hat{\mathbf{J}}(\mathbf{r}) \cdot [\hat{b}_{\mathbf{q}_s\lambda_s}, \hat{\mathbf{A}}(\mathbf{r})] \rangle \\ &= -i\omega_{\mathbf{q}_s} b_{\mathbf{q}_s\lambda_s} + \frac{ic\epsilon_{\mathbf{q}_s\lambda_s}}{\hbar\omega_{\mathbf{q}_s}} \int d\mathbf{r} e^{-i\mathbf{q}_s\cdot\mathbf{r}} \mathbf{e}_{\mathbf{q}_s\lambda_s} \cdot \mathbf{J}(\mathbf{r}, t). \end{aligned} \quad (48)$$

The second equality is obtained by an explicit computation of the commutator, in $\mathbf{J}(\mathbf{r}, t)$ as given in Eq. (16). Summary: It follows from Eqs. (46)–(48) and (16) that the heterodyne signal, introduced earlier in a fully quantum way and in terms of the field variables, can be expressed in terms of the expectation value of the gauge-invariant current density operator $\hat{\mathbf{J}}(\mathbf{r})$. This result is exact, and in particular can be considered as a starting point for calculating the radiative corrections to the signals. The latter, however, goes beyond the scope of this letter, and will be addressed elsewhere.

The heterodyne signal has the form

$$\begin{aligned} S_{\text{het}}(s) &= (\dot{E}_{\mathbf{q}_s\lambda_s}^*(t) \pm \dot{A}_{\mathbf{q}_s\lambda_s}^*(t)) \\ &\times \frac{ic\epsilon_{\mathbf{q}_s\lambda_s}}{\hbar\omega_{\mathbf{q}_s}} \int_{-\infty}^t dt \int d\mathbf{r} e^{i\omega_{\mathbf{q}_s} t - i\mathbf{q}_s\cdot\mathbf{r}} \mathbf{e}_{\mathbf{q}_s\lambda_s} \cdot \mathbf{J}(\mathbf{r}, t) \\ &+ (\bar{E}_{\mathbf{q}_s\lambda_s}^*(t) \pm \bar{A}_{\mathbf{q}_s\lambda_s}^*(t)) \\ &\times \frac{ic\epsilon_{\mathbf{q}_s\lambda_s}}{\hbar\omega_{\mathbf{q}_s}} \int d\mathbf{r} e^{i\omega_{\mathbf{q}_s} t - i\mathbf{q}_s\cdot\mathbf{r}} \mathbf{e}_{\mathbf{q}_s\lambda_s} \cdot \mathbf{J}(\mathbf{r}, t) + \text{c.c.} \end{aligned} \quad (49)$$

Under the semi-classical approximation, the interaction picture expansion of this expectation value in orders of incoming n vector fields is done in Eqs. (18)–(22). The heterodyne signal is thus given by the same response functions introduced for the energy exchange.

The above representation is gauge invariant, and it allows to reduce the summations to purely spatial values, and, finally provides a connection to the “standard” picture of optical response. Introducing formally the polarization \mathbf{P} by $\dot{\mathbf{P}} = -\mathbf{J}$, we can interpret Eq. (18), written in the $A_0 = 0$ gauge, and combined with Eq. (21) as an expansion of the system polarization in powers of the driving electric field. This is possible, despite the fact that there is no simple polarization operator, and the material system interacts with the driving field via the scalar and vector potentials, rather than the electric field. Introducing the polarization is not necessary in the present formalism and the comment was made to connect with the standard formalism.

V. CONCLUSIONS

We have shown that physical observables such as matter-field energy exchange and the heterodyne detected optical signal in Eq. (46) depend on expectation value of the gauge-invariant current density operator $\hat{\mathbf{J}}(\mathbf{r}, t; \mathbf{A})$. Note that $\sigma^+(\mathbf{r}, t)$ in Eqs. (24), (27), and (30) comes from the gauge-invariant current (Eq. (11)), whereas $\sigma^-(\mathbf{r}, t)$ comes from H_{int} in Eq. (4). We further presented a gauge-invariant non-local formal response

function in Liouville space to any order [Eq. (18)] and expanded it to first, second, and third orders [Eqs. (24), (27), and (30)], together with the corresponding Hilbert space representations [Eqs. (28), (28), and (31)]. The multipolar and minimal-coupling matter-field interaction Hamiltonians are consistent as shown in Eqs. (32)–(42). The complexity of the multipolar formalism grows rapidly for the non-linear optical response when using both electric and magnetic multipoles in nano-shaped lights,¹² which can be avoided in this formalism. The non-local response functions allow us to exactly calculate heterodyne detected optical signals in the presence of strong fields and non-uniform nano-optical fields.^{12,38} The formalism can be used to study non-adiabatic molecular current density dynamics.³⁹ Furthermore, it can be extended to cavity quantum electrodynamics (QED), or be used in understanding field angular momentum.⁴⁰

ACKNOWLEDGMENTS

The authors gratefully acknowledge the support of National Science Foundation (Grant Nos. CHE-1361516 and CHE-1111350) and the Chemical Sciences, Geosciences and Biosciences Division, Office of Basic Energy Sciences, Office of Science, U.S. Department of Energy through award #DE-FG02-04ER15571. The computational resources were provided by DOE.

APPENDIX: LAMB DISCREPANCY OF $(-i\omega_{\alpha 0}/\omega)^2$

We present the frequency domain calculation of linear non-local field/matter energy exchange which recovers the discrepancy of $(-i\omega_{\alpha 0}/\omega)^2$ in the linear absorption spectra when using “naive” minimal coupling Hamiltonian and dipole approximation.³⁷ In the dipole approximation, the field/matter couplings is given by^{1,2}

$$\hat{H}_{int}^d = -\hat{\boldsymbol{\mu}} \cdot \mathbf{E}(t), \quad (\text{A1})$$

where $\hat{\boldsymbol{\mu}}$ is dipole operator, $\mathbf{E}(t)$ is the electric field; in the dipole approximation, \mathbf{r} -dependence of the electric field is ignored. Thus, standard response theory gives us

$$\Delta W_d^{(1)}(t) = -i\hbar^{-1} \int_{-\infty}^t d\tau \dot{\mathbf{E}}(t) \cdot \langle \hat{\boldsymbol{\mu}}^+(t) \hat{\boldsymbol{\mu}}^-(\tau) \rangle \cdot \mathbf{E}(\tau), \quad (\text{A2})$$

or in frequency domain as

$$\begin{aligned} \Delta W_d^{(1)} &= \omega \int d\omega |\mathbf{E}(\omega)|^2 \mathcal{S}_d(\omega), \\ \mathcal{S}_d(\omega) &= -\hbar^{-1} \int_0^\infty d\tau \langle \boldsymbol{\mu}^+(\tau) \boldsymbol{\mu}^-(0) \rangle \exp(i\omega\tau). \end{aligned} \quad (\text{A3})$$

Expressing Eq. (A3) in sum over states and using $\mathbf{E}(\omega) = 2\pi E_0 \delta(\omega - \omega')$, we can write

$$\begin{aligned} \Delta W_d^{(1)} &= -2|E_0|^2 \omega \text{Im} \left\{ \left(\frac{n_0}{\hbar} \sum_{\alpha} [P(\alpha) - P(0)] \right. \right. \\ &\quad \left. \left. \times \frac{|\mu_{\alpha 0}|^2}{\omega - \omega_{\alpha 0} + i\eta_{\alpha 0}} \right) \right\}. \end{aligned} \quad (\text{A4})$$

Here, n_0 is the total number of particles per unit volume and $P(0)$, $P(\alpha)$ are the thermal population of state $|0\rangle$ $|\alpha\rangle$, respectively, and are defined in usual way. $\mu_{\alpha 0}$ and $\omega_{\alpha 0}$ are relevant

electric dipole elements and energy related to the electronic transition $|0\rangle \rightarrow |\alpha\rangle$.

A “naive” application of the minimal coupling Hamiltonian neglects the $\mathbf{A}^2(\mathbf{r}, t)$ since it is non-linear. This then gives

$$\Delta W_{mc}^{(1)}(t) = -i\hbar^{-1} \int_{-\infty}^t d\tau_1 \dot{\mathbf{A}}(t) \cdot \langle \hat{\mathbf{j}}^+(t) \hat{\mathbf{j}}^-(\tau_1) \rangle \cdot \mathbf{A}(\tau_1), \quad (\text{A5})$$

or in frequency domain as

$$\begin{aligned} \Delta W_{mc}^{(1)} &= \omega \int d\omega |\mathbf{A}(\omega)|^2 \mathcal{S}_{mc}(\omega), \\ \mathcal{S}_{mc}(\omega) &= -\hbar^{-1} \int_0^\infty d\tau \langle \hat{\mathbf{j}}^+(\tau) \hat{\mathbf{j}}^-(0) \rangle \exp(i\omega\tau) \\ &= \left(\frac{n_0}{\hbar} \sum_{\alpha} [P(\alpha) - P(0)] \frac{|\mathbf{j}_{\alpha 0}|^2}{\omega - \omega_{\alpha 0} + i\eta_{\alpha 0}} \right). \end{aligned} \quad (\text{A6})$$

The current density elements for given transition (say, $|0\rangle \rightarrow |\alpha\rangle$) $\mathbf{j}_{0\alpha} = \mathbf{j}_{\alpha 0}^* = \langle \alpha | \hat{\mathbf{j}} | 0 \rangle$ is related to *transition dipole* $\mu_{0\alpha}(t) = \mu_{0\alpha} \exp(-i\omega_{0\alpha} t)$ by

$$\mu_{0\alpha} = -i(\omega_{0\alpha})^{-1} \mathbf{j}_{0\alpha}, \quad (\text{A7})$$

since $\mathbf{E}(\omega) = i\omega \mathbf{A}(\omega)$, the first order matter/radiation energy exchange in *minimal coupling* thus becomes

$$\Delta W_{mc}^{(1)} = \left(\frac{-i\omega_{\alpha 0}}{\omega} \right)^2 \Delta W_d^{(1)}. \quad (\text{A8})$$

As can be seen from Eqs. (A3)–(A8), the matter/field energy exchange differ by a factor of $(-i\omega_{\alpha 0}/\omega)^2$ in the minimal coupling (ignoring $\mathbf{A}^2(\mathbf{r}, t)$) and dipole approximation. This factor adds asymmetry to the absorption peak and was resolved by Ref. 37 using tedious canonical transformation. This apparent discrepancy is caused by the “naive” use of the minimal coupling Hamiltonian rather than gauge invariant interaction as shown in main text under Sec. III.

¹S. Mukamel, *Principles of Nonlinear Optical Spectroscopy* (Oxford University Press, New York, 1995), Vol. 29.

²M. O. Scully, *Quantum Optics* (Cambridge University Press, 1997).

³C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Photons and Atoms: Introduction to Quantum Electrodynamics* (Wiley Professional, 1997).

⁴Y. Aharonov and D. Bohm, “Significance of electromagnetic potentials in the quantum theory,” *Phys. Rev.* **115**, 485 (1959).

⁵J. Rammer, *Quantum Transport Theory* (Westview Press, 2004).

⁶R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman Lectures on Physics, Desktop Edition Volume I* (Basic Books, 2013), Vol. 1.

⁷P. Bharadwaj, B. Deutsch, and L. Novotny, “Optical antennas,” *Adv. Opt. Photonics* **1**, 438–483 (2009).

⁸M. I. Stockman, “Nanoplasmonics: Past, present, and glimpse into future,” *Opt. Express* **19**, 22029–22106 (2011).

⁹M. I. Stockman, “Quantum nanoplasmonics,” *Photonics, Volume II: Scientific Foundations, Technology and Applications* (John Wiley & Sons, Inc., 2015), pp. 85–132.

¹⁰J. E. Donehue, E. Wertz, C. N. Talicska, and J. S. Biteen, “Plasmon-enhanced brightness and photostability from single fluorescent proteins coupled to gold nanorods,” *J. Phys. Chem. C* **118**, 15027–15035 (2014).

¹¹D. A. Kalashnikov, Z. Pan, A. I. Kuznetsov, and L. A. Krivitsky, “Quantum spectroscopy of plasmonic nanostructures,” *Phys. Rev. X* **4**, 011049 (2014).

¹²L. Novotny and B. Hecht, *Principles of Nano-Optics* (Cambridge University Press, 2012).

¹³F. Ding, E. B. Guidez, C. M. Aikens, and X. Li, “Quantum coherent plasmon in silver nanowires: A real-time TDDFT study,” *J. Chem. Phys.* **140**, 244705 (2014).

¹⁴R. M. Stöckle, Y. D. Suh, V. Deckert, and R. Zenobi, “Nanoscale chemical analysis by tip-enhanced Raman spectroscopy,” *Chem. Phys. Lett.* **318**, 131–136 (2000).

¹⁵E. Bailo and V. Deckert, “Tip-enhanced Raman scattering,” *Chem. Soc. Rev.* **37**, 921–930 (2008).

- ¹⁶M. L. Brongersma and P. G. Kik, *Surface Plasmon Nanophotonics* (Springer, 2007).
- ¹⁷S. Kühn, U. Håkanson, L. Rogobete, and V. Sandoghdar, "Enhancement of single-molecule fluorescence using a gold nanoparticle as an optical nanoantenna," *Phys. Rev. Lett.* **97**, 017402 (2006).
- ¹⁸R. Zhang, Y. Zhang, Z. C. Dong, S. Jiang, C. Zhang, L. G. Chen, L. Zhang, Y. Liao, J. Aizpurua, Y. Luo, J. L. Yang, and J. G. Hou, "Chemical mapping of a single molecule by plasmon-enhanced Raman scattering," *Nature* **498**, 82–86 (2013).
- ¹⁹L. V. Brown, X. Yang, K. Zhao, B. Zheng, P. Nordlander, and N. J. Halas, "Fan-shaped gold nanoantennas above reflective substrates for surface-enhanced infrared absorption (SEIRA)," *Nano Lett.* **15**, 1272 (2015).
- ²⁰A. Minovich, A. E. Klein, N. Janunts, T. Pertsch, D. N. Neshev, and Y. S. Kivshar, "Generation and near-field imaging of Airy surface plasmons," *Phys. Rev. Lett.* **107**, 116802 (2011).
- ²¹M. Aeschlimann, M. Bauer, D. Bayer, T. Brixner, F. J. G. de Abajo, W. Pfeiffer, M. Rohmer, C. Spindler, and F. Steeb, "Adaptive subwavelength control of nano-optical fields," *Nature* **446**, 301–304 (2007).
- ²²N. S. Abadeer, M. R. Brennan, W. L. Wilson, and C. J. Murphy, "Distance and plasmon wavelength dependent fluorescence of molecules bound to silica-coated gold nanorods," *ACS Nano* **8**, 8392–8406 (2014).
- ²³T. Taminiau, F. Stefani, F. Segerink, and N. Van Hulst, "Optical antennas direct single-molecule emission," *Nat. Photonics* **2**, 234–237 (2008).
- ²⁴T. Kudo and H. Ishihara, "Proposed nonlinear resonance laser technique for manipulating nanoparticles," *Phys. Rev. Lett.* **109**, 087402 (2012).
- ²⁵S. Palomba and L. Novotny, "Near-field imaging with a localized nonlinear light source," *Nano Lett.* **9**, 3801–3804 (2009).
- ²⁶V. Myroshnychenko, J. Rodríguez-Fernández, I. Pastoriza-Santos, A. M. Funston, C. Novo, P. Mulvaney, L. M. Liz-Marzán, and F. J. G. de Abajo, "Modelling the optical response of gold nanoparticles," *Chem. Soc. Rev.* **37**, 1792–1805 (2008).
- ²⁷K. Rzkazewski and R. Boyd, "Equivalence of interaction Hamiltonians in the electric dipole approximation," *J. Mod. Opt.* **51**, 1137–1147 (2004).
- ²⁸S. Tanaka, V. Chernyak, and S. Mukamel, "Time-resolved x-ray spectroscopies: Nonlinear response functions and Liouville-space pathways," *Phys. Rev. A* **63**, 063405 (2001).
- ²⁹A. Salam, *Molecular Quantum Electrodynamics: Long-Range Intermolecular Interactions* (John Wiley & Sons, 2010).
- ³⁰D. P. Craig and T. Thirunamachandran, *Molecular Quantum Electrodynamics: An Introduction to Radiation Molecular Interactions* (Academic Press, London, 1984).
- ³¹L. D. Romero, D. Andrews, and M. Babiker, "A quantum electrodynamics framework for the nonlinear optics of twisted beams," *J. Opt. B: Quantum Semiclassical Opt.* **4**, S66 (2002).
- ³²A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (Courier Corporation, 2003).
- ³³K. Cho, *Optical Response of Nanostructures: Microscopic Nonlocal Theory* (Springer Science & Business Media, 2003), Vol. 139.
- ³⁴C. A. Marx, U. Harbola, and S. Mukamel, "Nonlinear optical spectroscopy of single, few, and many molecules: Nonequilibrium Green's function QED approach," *Phys. Rev. A* **77**, 022110 (2008).
- ³⁵H. J. Werner, P. J. Knowles, G. Knizia, F. R. Manby, M. Schütz, P. Celani, T. Korona, R. Lindh, A. Mitrushenkov, G. Rauhut, K. R. Shamasundar, T. B. Adler, R. D. Amos, A. Bernhardsson, A. Berning, D. L. Cooper, M. J. O. Deegan *et al.*, MOLPRO, version 2012.1, a package of *ab initio* programs, 2012, see <http://www.molpro.net>.
- ³⁶E. Power and S. Zienau, *Philos. Trans. R. Soc., A* **251**, 427 (1959).
- ³⁷W. E. Lamb, Jr., R. R. Schlicher, and M. O. Scully, "Matter-field interaction in atomic physics and quantum optics," *Phys. Rev. A* **36**, 2763 (1987).
- ³⁸O. Keller, *Quantum Theory of Near-Field Electrodynamics* (Springer Science & Business Media, 2012), Vol. 2.
- ³⁹A. Mandal and K. L. C. Hunt, "Non-adiabatic current densities, transitions, and power absorbed by a molecule in a time-dependent electromagnetic field," *J. Chem. Phys.* **143**, 034102 (2015).
- ⁴⁰R. W. Heeres and V. Zwiller, "Subwavelength focusing of light with orbital angular momentum," *Nano Lett.* **14**, 4598–4601 (2014).